Water Parameters Tested By Mountain Regional Water D Showing levels tested for and violation information

Test Parameter #	Contaminant	Viola- tion (Y/N)	Units of Measure- ment	EPA Max. Cont- aminant Level Goal (MCLG)	EPA Max. Cont- aminant Level (MCL)	Likely Source(s) of Contamination	Required Health Effect Language (for Violation Notices)
A.	Microbiological and Particle Contaminant	s:					Outforward by the test of the
1	Total Coliform Bacteria	N	Cfu	0	note F	Naturally present in the environment	Coliforms are bacteria that are naturally present in the environment and are used as an indicator that other, potentially-harmful, bacteria may be present. Coliforms were found in more samples than allowed and this was a warning of potential problems. Fecal coliforms and E. coli are bacteria whose presence indicates that the water may be contaminated with human or animal wastes. Microbes in these wastes can cause short-term effects,
2	Fecal Coliform and E.coli	N	Cfu	0	note G	Human and animal fecal waste	such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a special health risk for infants, young children, some of the elderly, and people with severely compromised immune systems. Total organic carbon (TOC) has no health effects. However, total organic carbon provides a medium for the formation of disinfection byproducts. These byproducts include trihalomethanes
3	Total Organic Carbon (TOC)	N	ppm	n/a	π	Naturally present in the environment	(THMs) and haloacetic acids (HAAs). Drinking water containing these byproducts in excess of the MCL may lead to adverse health effects, liver or kidney problems, or nervous system effects, and may lead to an increased risk of getting cancer. Turbidity has no health effects. However, turbidity can interfere with disinfection and provide a medium for microbial growth. Turbidity may indicate the presence of disease-causing organisms.
4	Turbidity	N	NTU	n/a	TT	Soil runoff	These organisms include bacteria, viruses, and parasites that can cause symptoms such as nausea, cramps, diarrhea and associated headaches.
В.	Radioactive Contaminants:						
5	Beta/photon emitters	N	mrem/yr	0	4	Decay of natural and man-made deposits	Certain minerals are radioactive and may emit forms of radiation known as photons and beta radiation. Some people who drink water containing beta particle and photon radioactivity in excess of the MCL over many years may have an increased risk of getting cancer. Certain minerals are radioactive and may emit a form of radiation known as alpha radiation. Some people who drink water containing alpha emitters in excess of the MCL over many years may have
6	Alpha emitters	N	pCi/I	0	15	Erosion of natural deposits	an increased risk of getting cancer.
7	Combined Radium	N	pCi/I	0	5	Erosion of natural deposits	Some people who drink water containing radium 226 or 228 in excess of the MCL over many years may have an increased risk of getting cancer.
8	Uranium	N	pCi/l	0	30	Erosion of natural deposits	Some people who drink water containing uranium in excess of the MCL over many years may have an increased risk of getting cancer and kidney toxicity.
C.	Inorganic Contaminants:					Discharge from petroleum refineries; fire retardants;	Some people who drink water containing antimony well in excess of the MCL over many years could
9	Antimony	N	ppb	6	6	ceramics; electronics; solder	experience increases in blood cholesterol and decreases in blood sugar. Some people who drink water containing arsenic in excess of the MCL over many years could
10	Arsenic (see note H)	N	ppb	0	10	Erosion of natural deposits; Runoff from orchards; Runoff from glass and electronics production wastes	experience skin damage or problems with their circulatory system, and may have an increased risk of getting cancer.
11	Asbestos	N	MFL	7	7	Decay of asbestos cement water mains; Erosion of natural deposits	Some people who drink water containing asbestos in excess of the MCL over many years may have an increased risk of developing benign intestinal polyps.
12	Barium	N	ppm	2	2	Discharge of drilling wastes; Discharge from metal refineries; Erosion of natural deposits Discharge from metal refineries and coal-burning	Some people who drink water containing barium in excess of the MCL over many years could experience an increase in their blood pressure.
13	Beryllium	N	ppb	4	4	factories; Discharge from electrical, aerospace, and defense industries	Some people who drink water containing beryllium well in excess of the MCL over many years could develop intestinal lesions.
						Corrosion of galvanized pipes; Erosion of natural deposits; Dis-charge from metal refineries; Runoff from	Some people who drink water containing cadmium in excess of the MCL over many years could
14	Cadmium	N	ppb	5	5	waste batteries and paints Discharge from steel and pulp mills; Erosion of natural	experience kidney damage. Some people who use water containing chromium well in excess of the MCL over many years could
15	Chromium	N	ppb	100	100	deposits	experience allergic dermatitis.
							Copper is an essential nutrient, but some people who drink water containing copper in excess of the
16	Copper (see note C)	N	ppm	1.3	AL=1.3	Corrosion of household plumbing systems; Erosion of natural deposits; Leaching from wood preservatives	action level over a relatively short amount of time could experience gastrointestinal distress. Some people who drink water containing copper in excess of the action level over many years could suffer liver or kidney damage. People with Wilson's Disease should consult their personal doctor.
17	Cyanide	N	ppb	200	200	Discharge from steel/metal factories; Discharge from plastic and fertilizer factories	Some people who drink water containing cyanide well in excess of the MCL over many years could experience nerve damage or problems with their thyroid. Some people who drink water containing fluoride in excess of the MCL over many years could get
18	Fluoride	N	ppm	4	4	Erosion of natural deposits; Water additive which promotes strong teeth; Discharge from fertilizer and aluminum factories	bone disease, including pain and tenderness of the bones. Fluoride in drinking water at half the MCL or more may cause mottling of children's teeth, usually in children less than nine years old. Mottling, also known as dental fluorosis, may include brown staining and/or pitting of the teeth, and occurs only in developing teeth before they erupt from the gums. Infants and children who drink water containing lead in excess of the action level could experience
19	Lead (see note C)	N	ppb	0	AL=15	Corrosion of household plumbing systems; Erosion of natural deposits	delays in their physical or mental development. Children could show slight deficits in attention span and learning abilities. Adults who drink this water over many years could develop kidney problems or high blood pressure.

20	Mercury (inorganic)	N	ppb	2	2	and factories; Runoff from landfills; Runoff from cropland	Some people who drink water containing inorganic mercury well in excess of the MCL over many years could experience kidney damage. Infants below the age of six months who drink water containing nitrate in excess of the MCL could
21	Nitrate (as Nitrogen)	N	ppm	10	10	Runoff from fertilizer use; Leach-ing from septic tanks, sew-age; Erosion of natural deposits	become seriously ill and, if untreated, may die. Symptoms include shortness of breath and blue baby syndrome.
22	Nitrite (as Nitrogen)	N	ppm	1	1	Runoff from fertilizer use; Leach-ing from septic tanks, sew-age; Erosion of natural deposits	Infants below the age of six months who drink water containing nitrite in excess of the MCL could become seriously ill and, if untreated, may die. Symptoms include shortness of breath and blue baby syndrome. Selenium is an essential nutrient. However, some people who drink water containing selenium in
23	Selenium	N	ppb	50	50	Discharge from petroleum and metal refineries; Erosion	excess of the MCL over many years could experience hair or fingernail losses, numbness in fingers or toes, or problems with their circulation.
24	Thallium	N	ppb	0.5	2		Some people who drink water containing thallium in excess of the MCL over many years could experience hair loss, changes in their blood, or problems with their kidneys, intestines, or liver.
D.	Synthetic Organic Contaminants (including	y Pest	icides and Herbici	ides):			
25	2,4-D	N	ppb	70	70	Runoff from herbicide used on row crops	Some people who drink water containing the weed killer 2,4-D well in excess of the MCL over many years could experience problems with their kidneys, liver, or adrenal glands. Some people who drink water containing silvex in excess of the MCL over many years could
26	2,4,5-TP (Silvex)	N	ppb	50	50		experience liver problems. Some people who drink water containing high levels of acrylamide over a long period of time could
27	Acrylamide (see note D)	N		0	тт	Added to water during sewage/ wastewater treatment	have problems with their nervous system or blood, and may have an increased risk of getting cancer. Some people who drink water containing alachlor in excess of the MCL over many years could have
28	Alachlor	N	ppb	0	2	Runoff from herbicide used on row crops	problems with their eyes, liver, kidneys, or spleen, or experience anemia, and may have an increased risk of getting cancer.
29	Atrazine	N	ppb	3	3		Some people who drink water containing atrazine well in excess of the MCL over many years could experience problems with their cardiovascular system or reproductive difficulties.
30	Benzo(a)pyrene (PAH)	N	ppt	0	200	distribution lines	Some people who drink water containing benzo(a)pyrene in excess of the MCL over many years may experience reproductive difficulties and may have an increased risk of getting cancer.
31	Carbofuran	N	ppb	40	40	Leaching of soil fumigant used on rice and alfalfa	Some people who drink water containing carbofuran in excess of the MCL over many years could experience problems with their blood, or nervous or reproductive systems. Some people who drink water containing chlordane in excess of the MCL over many years could
32	Chlordane	N	ppb	0	2	Residue of banned termiticide	experience problems with their liver or nervous system, and may have an increased risk of getting cancer. Some people who drink water containing dalapon well in excess of the MCL over many years could
33	Dalapon	N	ppb	200	200	Runoff from herbicide used on rights of way	Some people who drink water containing dataport well in excess of the MCL over many years could experience minor kidney changes. Some people who drink water containing di (2-ethylhexyl) adipate well in excess of the MCL over
34	Di(2-ethylhexyl)adipate	N	ppb	400	400	Discharge from chemical factories	many years could experience general toxic effects or reproductive difficulties. Some people who drink water containing di (2-ethylhexyl) phthalate in excess of the MCL over many
35	Di(2-ethylhexyl)phthalate	N	ppb	0	6	Discharge from rubber and chem-ical factories	years may have problems with their liver, or experience reproductive difficulties, and may have an increased risk of getting cancer.
36	Dibromochloropropane	N	ppt	0	200	cotton, pineapples, and orchards	Some people who drink water containing DBCP in excess of the MCL over many years could experience reproductive difficulties and may have an increased risk of getting cancer.
37	Dinoseb	N	ppb	7	7	Runoff from herbicide used on soybeans and vegetables	
38	Diquat	N	ppb	20	20		Some people who drink water containing diquat in excess of the MCL over many years could get cataracts.
39	Dioxin[2,3,7,8-TCDD]	N	ppq	0	30		Some people who drink water containing dioxin in excess of the MCL over many years could experience reproductive difficulties and may have an increased risk of getting cancer.
40	Endothall	N		100	100	•	Some people who drink water containing endothall in excess of the MCL over many years could experience problems with their stomach or intestines.
41	Endrin	N	ppb	2	2		Some people who drink water containing endrin in excess of the MCL over many years could experience liver problems.
42	Epichlorohydrin (see note D)	N		0	TT	Discharge from industrial chemical factories; An impurity	Some people who drink water containing high levels of epichlorohydrin over a long period of time could experience stomach problems, and may have an increased risk of getting cancer.
							Some people who drink water containing ethylene dibromide in excess of the MCL over many years could experience problems with their liver, stomach, reproductive system, or kidneys, and may have
43	Ethylene dibromide	N	ppt	0	50	Discharge from petroleum refineries	an increased risk of getting cancer. Some people who drink water containing glyphosate in excess of the MCL over many years could
44	Glyphosate	N	ppb	700	700	Runoff from herbicide use	Some people who drink water containing heptachlor in excess of the MCL over many years could experience problems with their kidneys or reproductive difficulties. Some people who drink water containing heptachlor in excess of the MCL over many years could
45	Heptachlor	N	ppt	0	400	Residue of banned pesticide	Some people who drink water containing heptachlor epoxide in excess of the MCL over many years could experience liver damage and may have an increased risk of getting cancer. Some people who drink water containing heptachlor epoxide in excess of the MCL over many years
46	Heptachlorepoxide	N	ppt	0	200	Breakdown of heptachlor	could experience liver damage, and may have an increased risk of getting cancer. Some people who drink water containing hexachlorobenzene in excess of the MCL over many years
47	Hexachlorobenzene	N	ppb	0	1	Discharge from metal refineries and agricultural chemical factories	could experience problems with their liver or kidneys, or adverse reproductive effects, and may have an increased risk of getting cancer. Some people who drink water containing hexachlorocyclopentadiene well in excess of the MCL over
48	Hexachlorocyclopentadiene	N	ppb	50	50		many years could experience problems with their kidneys or stomach.

							_
49	Lindane	N	ppt	200	200	Runoff/leaching from insecticide used on cattle, lumber, gardens Runoff/leaching from insecticide used on fruits,	Son exp Son
50	Methoxychlor	N	ppb	40	40	vegetables, alfalfa, livestock	exp
51	Oxamyl [Vydate]	N	ppb	200	200	Runoff/leaching from insecticide used on apples, potatoes and tomatoes	Son exp Son
52	PCBs [Polychlorinatedbiphenyls]	N	ppt	0	500	Runoff from landfills; Discharge of waste chemicals	repr Son
53	Pentachlorophenol	N	ppb	0	1	Discharge from wood preserving factories	coul can Son
54	Picloram	N	ppb	500	500	Herbicide runoff	exp
55	Simazine	N	ppb	4	4	Herbicide runoff	exp
56 E.	Toxaphene Volatile Organic Contaminants:	N	ppb	0	3	Runoff/leaching from insecticide used on cotton and cattle	Son hav
57	Benzene	N	ppb	0	5	Discharge from factories; Leach-ing from gas storage tanks and landfills	exp can Son
58	Bromate	N	ppb	0	10	By-product of drinking water chlorination	an i
59	Carbon tetrachloride	N	ppb	0	5	Discharge from chemical plants and other industrial activities	Son coul Son irrita
60	Chloramines	N	ppm	MRDLG=4	MRDL=4	Water additive used to control microbes	Son
61	Chlorine	N	ppm	MRDLG=4	MRDL=4	Water additive used to control microbes	exc
62	Chlorite	N	ppm	0.8	1	By-product of drinking water chlorination	Son exp drin Son cou
63	Chlorine dioxide	N	ppb	MRDLG=800	MRDL=800	Water additive used to control microbes	who
64	Chlorobenzene	N	ppb	100	100	Discharge from chemical and agricultural chemical factories	Son
65	o-Dichlorobenzene	N	ppb	600	600	Discharge from industrial chemical factories	Son yea
66	p-Dichlorobenzene	N	ppb	75	75	Discharge from industrial chemical factories	Son cou Son
67	1,2 - Dichloroethane	N	ppb	0	5	Discharge from industrial chemical factories	may Son
68	1,1 - Dichloroethylene	N	ppb	7	7	Discharge from industrial chemical factories	yea Son
69	cis-1,2-ichloroethylene	N	ppb	70	70	Discharge from industrial chemical factories	yea Son
70	trans - 1,2 -Dichloroethylene	N	ppb	100	100	Discharge from industrial chemical factories	mar Son
71	Dichloromethane	N	ppb	0	5	Discharge from pharmaceutical and chemical factories	cou
72	1,2-Dichloropropane	N	ppb	0	5	Discharge from industrial chemical factories	may Son
73	Ethylbenzene	N	ppb	700	700	Discharge from petroleum refineries	cou
74	Haloacetic Acids [HAA] (see Note B)	N	ppb	n/a	60	By-product of drinking water disinfection Discharge from rubber and plastic factories; Leaching	may
75	Styrene	N	ppb	100	100	from land-fills	hav
76	Tetrachloroethylene	N	ppb	0	5	Discharge from factories and dry cleaners	cou
77	1,2,4 -Trichlorobenzene	N	ppb	70	70	Discharge from textile-finishing factories	mar

Some people who drink water containing lindane in excess of the MCL over many years could experience problems with their kidneys or liver.

Some people who drink water containing methoxychlor in excess of the MCL over many years could experience reproductive difficulties.

Some people who drink water containing oxamyl in excess of the MCL over many years could experience slight nervous system effects.

Some people who drink water containing PCBs in excess of the MCL over many years could experience changes in their skin, problems with their thymus gland, immune deficiencies, or reproductive or nervous system difficulties, and may have an increased risk of getting cancer. Some people who drink water containing pentachlorophenol in excess of the MCL over many years could experience problems with their liver or kidneys, and may have an increased risk of getting cancer.

Some people who drink water containing picloram in excess of the MCL over many years could experience problems with their liver.

Some people who drink water containing simazine in excess of the MCL over many years could experience problems with their blood.

Some people who drink water containing toxaphene in excess of the MCL over many years could have problems with their kidneys, liver, or thyroid, and may have an increased risk of getting cancer.

Some people who drink water containing benzene in excess of the MCL over many years could experience anemia or a decrease in blood platelets, and may have an increased risk of getting cancer.

Some people who drink water containing bromate in excess of the MCL over many years may have an increased risk of getting cancer.

Some people who drink water containing carbon tetrachloride in excess of the MCL over many years could experience problems with their liver and may have an increased risk of getting cancer. Some people who use water containing chloramines well in excess of the MRDL could experience irritating effects to their eyes and nose. Some people who drink water containing chloramines well in excess of the MRDL could experience stomach discomfort or anemia.

Some people who use water containing chlorine well in excess of the MRDL could experience irritating effects to their eyes and nose. Some people who drink water containing chlorine well in excess of the MRDL could experience stomach discomfort.

Some infants and young children who drink water containing chlorite in excess of the MCL could experience nervous system effects. Similar effects may occur in fetuses of pregnant women who drink water containing chlorite in excess of the MCL. Some people may experience anemia. Some infants and young children who drink water containing chlorine dioxide in excess of the MRDL could experience nervous system effects. Similar effects may occur in fetuses of pregnant women who drink water containing chlorine dioxide in excess of the MRDL. Some people may experience anemia.

Some people who drink water containing chlorobenzene in excess of the MCL over many years could experience problems with their liver or kidneys.

Some people who drink water containing o-dichlorobenzene well in excess of the MCL over many years could experience problems with their liver, kidneys, or circulatory systems.

Some people who drink water containing p-dichlorobenzene in excess of the MCL over many years could experience anemia, damage to their liver, kidneys, or spleen, or changes in their blood. Some people who drink water containing 1,2-dichloroethane in excess of the MCL over many years may have an increased risk of getting cancer.

Some people who drink water containing 1,1-dichloroethylene in excess of the MCL over many vears could experience problems with their liver.

Some people who drink water containing cis-1,2-dichloroethylene in excess of the MCL over many years could experience problems with their liver.

Some people who drink water containing trans-1,2-dichloroethylene well in excess of the MCL over many years could experience problems with their liver.

Some people who drink water containing dichloromethane in excess of the MCL over many years could have liver problems and may have an increased risk of getting cancer.

Some people who drink water containing 1,2-dichloropropane in excess of the MCL over many years

may have an increased risk of getting cancer.

Some people who drink water containing ethylbenzene well in excess of the MCL over many years

could experience problems with their liver or kidneys.

Some people who drink water containing haloacetic acids in excess of the MCL over many years

some people who drink water containing haloacetic acids in excess of the MCL over many years may have an increased risk of getting cancer.

Some people who drink water containing styrene well in excess of the MCL over many years could have problems with their liver, kidneys, or circulatory system.

Some people who drink water containing tetrachloroethylene in excess of the MCL over many years could have problems with their liver, and may have an increased risk of getting cancer. Some people who drink water containing 1,2,4-trichlorobenzene well in excess of the MCL over many years could experience changes in their adrenal glands.

						50 1 4 4 1 1 1 1 1 1 1 1 1
78	1,1,1 - Trichloroethane	N	ppb	200	200	Discharge from metal degreasing sites and other factories
79	1,1,2 -Trichloroethane	N	ppb	3	5	Discharge from industrial chemical factories
80	Trichloroethylene	N	ppb	0	5	Discharge from metal degreasing sites and other factories
81	TTHM [Total trihalomethanes] (see note B)	N	ppb	n/a	100/80	By-product of drinking water chlorination
82	Toluene	N	ppm	1	1	Discharge from petroleum factories Leaching from PVC piping; Discharge from plast
83	Vinyl Chloride	N	ppb	0	2	factories
						Discharge from petroleum factories; Discharge fr
84	Xylenes	N	ppm	10	10	chemical factories
F.	Secondary & Unregulated Standards:					
85	Iron	N	ppb		300	Erosion of natural deposits
86	Manganese	N	ppb		50	Erosion of natural deposits
87	Zinc	N	ppm		5	Erosion of natural deposits
88	Chloride	N	ppm		250	Erosion of natural deposits and road salting
89	Phosphorus	N	ppm		UR	Erosion of natural deposits
90	Sulfate	N	ppm		250	Erosion of natural deposits
91	Total Dissolved Solids or (TDS)	N	ppm		500	Erosion of natural deposits
92	Calcium	N	ppm		UR	Erosion of natural deposits
93	Hardness as CaCO3	N	ppm		UR	Erosion of natural deposits
94	Alkalinity - Total	N	ppm		UR	Erosion of natural deposits
95	Magnesium	N	ppm		UR	Erosion of natural deposits
96	Potassium	N	ppm		UR	Erosion of natural deposits
97	Sodium	N	ppm		UR	Erosion of natural deposits and road salting
98	Silica (as SiO2)	N	ppm		UR	Erosion of natural deposits
99	Aluminum	N	ppm		0.2	Erosion of natural deposits
100	Ammonia	N	ppm		UR	Sewage/Fertilizer
101	Boron	N	ppm		UR	Erosion of natural deposits
102	Hexavalant Chromium	N	ppm		UR	Erosion of natural deposits
103	Bicarbonate	N	ppm		UR	Erosion of natural deposits
104	Carbonate	N	ppm		UR	Erosion of natural deposits
105	Carbon Dioxide	N	ppm		UR	Erosion of natural deposits
106	Hydroxide	N	ppm		UR	Erosion of natural deposits
107	Specific Conductivity	N	umhos/cm		UR	Erosion of natural deposits
108	Silver	N	ppm		0.1 0.5	Erosion of natural deposits
109	Foaming Agents	N	ppm		0.5 Non-Corrosive	Erosion of natural deposits
110	Corosivity	N				Erosion of natural deposits
111	Color	N	color units		15	Erosion of natural deposits
112	Odor (see note E)	N	TON		3 6.5-8.5	Erosion of natural deposits
113	pH (in units)	N	pH units		0.5-8.5	Erosion of natural deposits

Some people who drink water containing 1,1,1-trichloroethane in excess of the MCL over many years could experience problems with their liver, nervous system, or circulatory system. Some people who drink water containing 1,1,2-trichloroethane well in excess of the MCL over many years could have problems with their liver, kidneys, or immune systems.

Some people who drink water containing trichloroethylene in excess of the MCL over many years could experience problems with their liver and may have an increased risk of getting cancer. Some people who drink water containing trihalomethanes in excess of the MCL over many years may experience problems with their liver, kidneys, or central nervous systems, and may have an increased risk of getting cancer.

Some people who drink water containing toluene well in excess of the MCL over many years could have problems with their nervous system, kidneys, or liver.

Some people who drink water containing vinyl chloride in excess of the MCL over many years may have an increased risk of getting cancer.

Some people who drink water containing xylenes in excess of the MCL over many years could experience damage to their nervous system.